

## Review

### \*Corresponding author

**Gustavo Santos Masson, PhD**  
Department of Physiology and  
Biophysics, Institute of Biomedical  
Sciences, University of Sao Paulo  
Av. Prof. Lineu Prestes, 1524 - ICB-I  
05508-000 Butantan - Sao Paulo/SP  
Brazil  
E-mail: [gsmasson@icb.usp.br](mailto:gsmasson@icb.usp.br)

Volume 2 : Issue 2

Article Ref. #: 1000HROJ2111

### Article History

Received: April 10<sup>th</sup>, 2015

Accepted: April 22<sup>nd</sup>, 2015

Published: April 23<sup>rd</sup>, 2015

### Citation

Masson GS, Michelini LC. Autonomic dysfunction, sympathetic hyperactivity and the development of end-organ damage in hypertension: Multiple benefits of exercise training. *Heart Res Open J.* 2015; 2(2): 60-69. doi: [10.17140/HROJ-2-111](http://dx.doi.org/10.17140/HROJ-2-111)

### Copyright

©2015 Masson GS. This is an open access article distributed under the Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

# Autonomic Dysfunction, Sympathetic Hyperactivity and the Development of End-Organ Damage in Hypertension: Multiple Benefits of Exercise Training

**Gustavo Santos Masson\* and Lisete Compagno Michelini**

*Department of Physiology and Biophysics, Biomedical Sciences Institute, University of Sao Paulo, Sao Paulo, SP, Brazil*

### ABSTRACT

Autonomic dysfunction is closely related to the development of hypertension, which is characterized by increased sympathetic activity, decreased vagal tonus and baroreflex dysfunction. The hypertension-induced maladaptive changes progressively lead to heart failure, myocardial infarction and stroke. Hypertrophic remodeling of brain arterioles, chemoreceptors activation, blood-brain barrier abnormalities, oxidative stress and pro-inflammatory cytokines production in autonomic brain areas increase neuronal activity and sympathetic outflow. These responses, together with increased baroreflex dysfunction-induced pressure variability, renin-angiotensin system hyperactivation and capillary rarefaction, increase blood pressure levels and act as a positive feedback mechanism to perpetuate hypertension and development of end-organ damage. Exercise training, a non-pharmacological tool, has been used as an adjuvant therapy to treat hypertension. Our recent data showed that moderate aerobic training in adult SHR completely normalizes oxidative stress and inflammation in autonomic brain areas involved in cardiovascular control and promptly corrects baroreflex dysfunction and increases cardiac vagal activity. The early (2-weeks) training-induced beneficial responses improve autonomic control even in the persistence of hypertension, since a partial reduction of pressure levels was observed after 8 weeks of exercise training, which was related to reversion of arteriolar hypertrophic remodeling and consequent decrease of peripheral vascular resistance.

**KEYWORDS:** Hypertension; Oxidative stress; Inflammation; Central nervous system; Baroreflex; Aerobic exercise training.

**ABBREVIATIONS:** ANG II: Angiotensin II; HMGB1: High-mobility group protein 1; HPLC: High Performance Liquid Chromatography; IL-1 $\beta$ : Interleukin 1 beta; MAPK: Mitogen-activated protein kinases; NADPH oxidase: Nicotinamide Adenine Dinucleotide Phosphate-oxidase; NF-K $\kappa$ : Nuclear factor kappa-light-chain-enhancer of activated B cells; NOS: Nitric Oxide Synthase; NTS: tractussolitarii nucleus; PKC: Protein Kinase C; PVN: hypothalamic paraventricular nucleus; RAS: Renin-angiotensin system; RVLM: Rostrolateral medulla; SFO: Sub-fornical organ; SHR: Spontaneously Hypertensive Rat; TGF- $\beta$ : Transforming Growth Factor beta; TNF- $\alpha$ : Tumoral Necrosis Factor alpha.

### INTRODUCTION

Central nervous system continuously monitors cardiovascular function through different types of receptors, which allow immediate and chronic hemodynamic adjustments evoked by endogenous and environmental stimuli. These adjustments are generated by the activation and/or inhibition of autonomic brain circuitry in the brain stem and the hypothalamus. Baroreceptors, chemoreceptors and cardiopulmonary receptors encode arterial pressure, partial pres-

sure of blood gases and cardiac filling changes in frequency of action potentials of respective afferents. This information is integrated in the central nervous system, which modifies vagal and sympathetic activity to heart and vasculature and, consequently, corrects the initial alterations and keeps arterial pressure, PO<sub>2</sub> and PCO<sub>2</sub> and stroke volume at relatively constant levels. The imbalance between activation/inhibition of autonomic control areas promotes chronic adaptations in cardiovascular effectors that contribute to the establishment of arterial hypertension and the development of end-organ injuries.<sup>1</sup> The mechanisms underlying the establishment of autonomic dysfunction, sympathetic hyperactivity and the appearance of hypertensive dysfunctions will be briefly reviewed. In addition, we will address the benefits of aerobic training in an experimental model of primary hypertension, the Spontaneously Hypertensive Rat (SHR).

#### AUTONOMIC DYSFUNCTION IN ARTERIAL HYPERTENSION

Although hypertension *per se* does not induce death, the pathological mechanisms associated with chronic arterial pressure elevation contribute to the development of end-organ injuries, as cardiac hypertrophy, stroke and glomerular sclerosis, which increase cardiovascular mortality. Since arterial hypertension is a multifactorial syndrome, many mechanisms overlapped to produce a positive feedback that facilitates cardiovascular dysfunction. Among these mechanisms, autonomic dysfunction seems to be a key factor in the pathophysiology of primary hypertension<sup>2</sup> and consists an important pharmacological target for blood pressure control and for the reduction of morbimortality.<sup>3</sup>

Autonomic dysfunction is characterized by the increased sympathetic nerve activity, decreased vagal nerve activity and abnormal reflex control of cardiovascular function, which are mediated by baroreceptors, chemoreceptors and cardiopulmonary receptors. Causative relation between autonomic dysfunction and elevated arterial pressure was primarily suggested in the 80's by Minami and co-authors<sup>4</sup> that identified reduced reflex bradycardia on juvenile SHR with normal arterial pressure. This finding indicated that autonomic dysfunction preceded the arterial pressure rise.

Many hypothesis have been proposed to explain the central nervous system abnormalities conditioning autonomic dysfunction in primary hypertension: baroreflex dysfunction itself, increased chemoreflex activation induced by reduced cerebral blood flow associated with arteriolar remodeling,<sup>5</sup> activation of peripheral chemosensitive cells,<sup>6,7</sup> increased brain Renin-angiotensin system (RAS),<sup>8,9</sup> as well increased blood-brain barrier permeability.<sup>10</sup> An elegant series of experiments developed by Julian Paton's group at the Bristol University identified the contribution of brain arteries/arterioles hypertrophic remodeling in the establishment of sympathetic hyperactivity. It was observed that neonates SHR had greater arterial wall thickness and elevated vertebral artery wall/lumen ratio with increased vascular resistance in the vertebro-basilar circuit, which determined reduced blood flow and, consequently, the rise of sympathetic

activity and peripheral vasoconstriction.<sup>5,11</sup>

Although not well characterized as the role of brain vasculature hypertrophic remodeling, it was also suggested that activation of chemosensitive afferents by pro-inflammatory cytokines contributed to the activation of sympathetic activity.<sup>6,12</sup> Indeed, several studies demonstrated a direct correlation between plasma pro-inflammatory cytokines, increased arterial pressure<sup>13</sup> and autonomic dysfunction.<sup>14</sup> Mkrtchian and co-authors<sup>15</sup> identified, in humans, some inflammatory mediators (toll-like receptor 1 and 4, HMGB1, TNF- $\alpha$  and IL-1 $\beta$  receptors and transcription factor NF- $\kappa$ B) within the carotid bodies, where chemosensitive cells are located. Hyperactivity of chemosensitive afferents was also observed in young SHR, since denervation of the carotid body caused reduction of arterial pressure, sympathetic activity and macrophages infiltration in the smooth muscle tissue.<sup>7</sup> Together, these findings suggested that pro-inflammatory cytokines-induced activation of chemoreceptors contributed to establishment of autonomic dysfunction and, consequently, of hypertension.

Another contributor factor to the genesis of sympathetic hyperactivity is ANG II-induced AT<sub>1</sub> receptors activation in brain areas without blood-brain barrier, as the Sub-fornical organ (SFO), which modulates other pre-autonomic areas, such as the hypothalamic Paraventricular nucleus (PVN) and the Rostrolateral medulla (RVL). RAS inhibition in these areas abolishes arterial pressure increase, renal sympathetic hyperactivity, baroreflex dysfunction and attenuates the dipsogenic response. RAS inhibition also corrects the expression of pro-inflammatory cytokines and AT<sub>1</sub> receptors and reduces reactive oxygen species content in ANG II-dependent hypertension.<sup>8,9,16</sup> renovascular hypertension<sup>17</sup> and primary hypertension.<sup>18</sup> These data show that increased AT<sub>1</sub> receptor activation, *via* reactive oxygen species and pro-inflammatory cytokines, contribute to the development of autonomic dysfunction, sympathetic hyperactivity and hypertension. Activation of AT<sub>1</sub> receptors also disrupts blood brain barrier and facilitates the migration of monocytes and T cells into brain, thus contributing to local inflammation and to a further increase in sympathetic activity and arterial pressure.<sup>10</sup> It was demonstrated that hematopoietic cells bind to junctional adhesion molecule-1 to enter into the neural tissue, where they behaved as resident macrophages (activated microglia), acting as another important source for pro-inflammatory cytokines.<sup>19</sup>

#### MOLECULAR MECHANISMS OF SYMPATHETIC HYPERACTIVITY: THE IMPORTANCE OF REACTIVE OXYGEN SPECIES AND PRO-INFLAMMATORY CYTOKINES

Several mechanisms that induce autonomic dysfunction exhibit common factors: increased reactive oxygen species and pro-inflammatory cytokines in autonomic brain areas as the SFO, PVN, RVL and tractus solitarius nucleus (NTS). The main intracellular signaling pathways in neurons and glia are activated by ANG II *via* AT<sub>1</sub> receptor, which activates NADPH oxidase through the Protein Kinase C (PKC) with the subsequent

release of superoxide.<sup>20-23</sup> Increased superoxide production activates redox-sensitive pathways, as the Mitogen-Activated Protein Kinases (MAPK) that stimulate nuclear transcription factors (NF- $\kappa$ B and AP-1), thus increasing the gene expression of pro-inflammatory cytokines, as well as subunits of the NADPH oxidase and others RAS components.<sup>22,24,25</sup> Therefore, AT<sub>1</sub> receptors activation, oxidative stress and inflammation constitute important positive feedback mechanisms in autonomic dysfunction and sympathetic hyperactivity.

Besides the regulation of nuclear factors, reactive oxygen species directly increase sympathetic neuronal activity. In ANG II-dependent hypertension, Robin Davisson and Constance Iadecola's groups, at the Cornell University, identified that NADPH oxidase-induced superoxide production increases the calcium influx and neuronal activation *via* glutamatergic NMDA receptors activation.<sup>21,23,26</sup> As calcium influx was abolished by antioxidant agents or nitric oxide donors,<sup>26</sup> the authors proposed that increased reactive oxygen species production decreased nitric oxide bioavailability and, consequently, NMDA receptor NR1 subunit nitrosylation,<sup>27</sup> thus increasing both neuronal activity and sympathetic activity. Several studies in ANG II-dependent hypertension,<sup>28-30</sup> renovascular hypertension<sup>31-33</sup> and primary hypertension,<sup>34-36</sup> identified that oxidative stress attenuation inhibits both arterial pressure and renal sympathetic activity elevation, decreases tissue RAS, pro-inflammatory cytokines and NADPH oxidase expression and reduces NF- $\kappa$ B e AP-1 transcriptional activity. These studies confirm the functional role of reactive oxygen species in autonomic dysfunction, sympathetic hyperactivity and, subsequently, in the development/establishment of arterial hypertension. In addition to sympathetic hyperactivity, reactive oxygen species also mediates lower parasympathetic activity and reduced baroreflex sensitivity after ANG II administration into the NTS.<sup>37</sup>

Tissue inflammation is another mechanism related to the establishment of autonomic dysfunction. It was described that healthy rats that received pro-inflammatory cytokines (TNF- $\alpha$ , IL-1 $\beta$  and IL-6) into the PVN and NTS or through intracarotid injection,<sup>38-40</sup> exhibited autonomic dysfunction, increased renal sympathetic activity and elevated arterial pressure. In addition to these acute effects of cytokines, Joseph Francis's group, at Louisiana State University, demonstrated that chronic blockade of pro-inflammatory cytokines expression into autonomic control areas attenuated cardiac hypertrophy, renal sympathetic hyperactivity and the elevation of arterial pressure in ANG II-induced hypertension.<sup>8,25</sup>

#### SYMPATHETIC HYPERACTIVITY AND END-ORGAN DAMAGE IN HYPERTENSION

Adrenergic hyperactivity in hypertension was originally described by Amann, et al.<sup>41</sup> which identified increased forearm blood flow after  $\alpha$ -adrenergic receptors blockade. The interaction between norepinephrine and  $\alpha$ -adrenergic receptors in smooth muscle cells induced vasoconstriction through

increased calcium influx and decreased nitric oxide bioavailability that elevated the peripheral vascular resistance and the arterial pressure.<sup>42</sup> Increased vascular sympathetic activity produced hypertrophic remodeling of arteries and arterioles,<sup>43</sup> since  $\alpha$ -adrenergic stimulation increases the expression of adhesion molecules, leukocyte migration, activation of NADPH oxidase with reactive oxygen species formation and MAPKs stimulation, which drive hypertrophic vascular effects.<sup>44</sup>

In our group, neuronal recordings, dosage of norepinephrine content in vessels (HPLC) and tyrosine hydroxylase immunoreactivity in different tissues were used to demonstrate that increased sympathetic vasomotor tonus in SHR is not homogenous, since we observed increased sympathetic activity in cardiac and renal, but not in skeletal muscle arterioles.<sup>45</sup> Different sympathetic activation patterns, as increased renal and cardiac sympathetic nerve activity and unchanged lumbar nerve activity,<sup>46</sup> were described in others hypertensive models, defining a differential *sympathetic signature*. Interestingly the *sympathetic signature* of the SHR coincides with the sympathetic activation pattern observed in hypertensive patients.

Besides vascular hypertrophy, increased peripheral vascular resistance and elevated arterial pressure, sympathetic hyperactivity contributes significantly to the development of end-organ injuries. In the myocardium, the higher metabolic demand caused by sympathetic hyperactivity (increased vascular resistance and elevated heart rate) is an important factor to determine ventricular hypertrophy. Neurohormonal direct effects of sympathetic hyperactivity were described in the myocardium. It was observed that cardiac  $\alpha$ - and  $\beta$ -adrenergic activation induced cardiac hypertrophy, augmented matrix metalloproteinase-2 activity, increased the expression of TGF- $\beta$  and the synthesis of collagen I and III, in addition to intensify the production of reactive oxygen species and the infiltration of hematopoietic mononuclear cells.<sup>47-49</sup> Accordingly, Schlaich and co-authors<sup>50</sup> identified in hypertensive patients a direct correlation between cardiac norepinephrine spillover and cardiac hypertrophy.

Similar to cardiac and vascular tissues, adrenergic hyperactivation, by modifying sodium/water reabsorption<sup>51</sup> and renin secretion,<sup>52</sup> causes abnormalities in the renal function. Graham and co-authors<sup>53</sup> described increased renal  $\alpha$ -adrenergic receptor concentration in the SHR. It was also demonstrated that subpressor doses of  $\alpha$ - and  $\beta$ -adrenergic blockers decreased glomerular sclerosis and urinary albumin excretion in partial nephrectomized rats.<sup>54</sup> Besides these direct effects, increased renal sympathetic activity determined tissue and plasma RAS activation through juxtaglomerular cells that release renin. Acting in AT<sub>1</sub> receptors, ANG II induced reactive oxygen species and pro-inflammatory cytokines production that enabled renal remodeling and injury, as the glomerular sclerosis.<sup>55</sup> Renin also interacts with pro-renin receptors in several tissues being associated with additional activation of oxidative and inflammatory signaling pathways, and, consequently, with the worsening of end-organ injuries.<sup>56,57</sup> ANG II systemic and local effects in the cardiac<sup>58</sup>

and renal<sup>59</sup> tissues add to brain RAS effects,<sup>8,9,16,17,18,21,28,29</sup> to facilitate sympathetic hyperactivity and to amplify the deleterious effects of hypertension.

Other mechanism closely related to end-organ injuries is the reduction of baroreflex sensitivity, which strengthens sympathetic hyperactivity. In this sense, Nosaka and co-authors<sup>60</sup> demonstrated that RVLM activation inhibited baroreceptors activation-induced reflex bradycardia, thus aggravating baroreflex dysfunction that is considered an independent prognostic marker in hypertension.<sup>61</sup> Baroreflex dysfunction decreases the ability of arterial baroreceptors to promptly correct venous return, heart rate, ventricular contractility and peripheral vascular resistance changes, which aggravate arterial pressure oscillations. High pressure variability, which increases hydrostatic pressure oscillations in the capillaries, expose tissues to brief periods of hyperperfusion or hypoperfusion interfering with normal tissue oxygenation. Hypoxia or partial oxygen pressure fall is a strong stimulus to drive endothelial cell injury and capillary apoptosis, which determine extensive capillary rarefaction and the consequent development of lesions in the various target organs.<sup>62,63</sup>


In summary, as illustrated in Figure 1, hypertension is associated to brain RAS hyperactivation, increased reactive oxygen species and pro-inflammatory cytokines in the autonomic brain areas that determine baroreflex dysfunction, decreased vagal cardiac activity and increased sympathetic activity to cardiac, vascular and renal tissues and increased pressure variability. These effects cause sympatho-vagal imbalance in the heart, activation of systemic and local RAS in peripheral tissues, vascular and tissue hypertrophic remodeling and end-organ damage.

The deleterious adaptive responses potentiate the development/maintenance of hypertension and constitute a positive feedback mechanism to perpetuate the hypertensive disease.

#### AEROBIC TRAINING: AUTONOMIC BENEFITS TO HYPERTENSIVE INDIVIDUALS

Moderate intensity exercise training is one of the most important non-pharmacological strategies to decrease arterial pressure in hypertensive patients. A recent meta-analysis demonstrated that aerobic training decreases systolic and diastolic arterial pressure by 8 and 5 mm Hg, respectively.<sup>65</sup> Other works also recognized that aerobic training corrects autonomic dysfunction associated with hypertension, which contributes to the reduction of end-organ damage and cardiovascular mortality.<sup>66-68</sup>

Clinical<sup>69</sup> and experimental<sup>70-74</sup> studies have indicated that aerobic training is extremely efficient to revert autonomic dysfunction and to attenuate sympathetic hyperactivity, to normalize arterioles wall/lumen ratio and to decrease peripheral vascular resistance. On never-treated hypertensive patients, Latorza and co-authors<sup>69</sup> identified that aerobic training normalized reflex control of heart rate and decreased sympathetic activity. On SHR, several studies from our group demonstrated that aerobic training decreases sympathetic neuronal excitability into the PVN<sup>75</sup> and increased PVN density of oxytocin neurons and the density of oxytocinergic projections from PVN to NTS-DMV complex,<sup>76</sup> whose activation increase both baroreflex sensitivity and cardiac vagal activity and decrease peripheral vasomotor sympathetic activity. It was also demonstrated that the training-induced adaptive responses are significantly correlated with de-



**Figure1:** Positive feedback mechanisms perpetuating hypertension. Increased RAS activation, reactive oxygen species and pro-inflammatory cytokines in autonomic brain areas of hypertensive individuals augments neuronal activity and sympathetic outflow and decreases both vagal activity and baroreflex sensitivity. The autonomic dysfunction increases arterial pressure variability and facilitates peripheral RAS activity and vascular and tissue remodeling, all these effects contributing to end-organ damage and to the maintenance of hypertension. NTS: Nucleus of the solitary tract; PVN: Paraventricular nucleus of the hypothalamus; RAS: Renin-angiotensin system; RVLM: Rostrolateral medulla. Modified from Masson & Michelini, 2014.<sup>64</sup>

creased resting heart rate and exercise tachycardia.<sup>77,78</sup>

In a recent study,<sup>70</sup> we identified the sequential changes of autonomic and cardiovascular adaptations induced by aerobic training in SHR (Figure 2). Only 2-weeks of aerobic training were enough to normalize cardiac vagal activity and baroreflex sensitivity in the SHR. There was, at the same experimental time, a complete normalization of the oxidative stress and inflammatory profile into the PVN, which are the most prominent molecular mechanisms to normalize baroreflex dysfunction and sympathetic hyperactivity. It is important to note that these benefits were observed even in the persistence of hypertension, since a partial decrease of arterial pressure (~9%) was identified only after 8-weeks of aerobic training (Figure 2).<sup>70</sup> The temporal coincidence between the reversion of pro-inflammatory profile and oxidative stress in the PVN and the correction of autonomic dysfunction,<sup>70</sup> associated with previous findings proving the relationship among oxidative stress,<sup>8,31,34</sup> inflammation,<sup>8,25</sup> baroreflex dysfunction and sympathetic hyperactivity,<sup>39,40</sup> suggest a cause-effect relationship. In other words, aerobic training normalizes baroreflex function and decreases sympathetic hyperactivity through the correction of pro-inflammatory profile and oxidative stress within the PVN, an important autonomic area in cardiovascular control. This statement is corroborated by another study from our group that observed that aerobic training decreases neuronal excitability in the PVN of the SHR.<sup>75</sup>


It is important to note the temporal dissociation between a prompt normalization of both baroreflex function and cardiac vagal activity (significant at 2-weeks) and arterial pressure fall, which appeared only after 8-weeks of exercise training.<sup>70</sup> This dissociation suggests that training-induced normalization of baroreflex function is independent of pressure reduction, but may

contribute to the subsequent pressure fall through the decrease of sympathetic vasomotor activity and the regression of vascular hypertrophy. Indeed, the contribution of increased baroreflex sensitivity to pressure fall was identified in clinical studies<sup>79,80</sup> that used chronic baroreflex stimulation to control refractory hypertension. The importance of augmented baroreflex function for a better cardiovascular control after aerobic training was also proved by other studies<sup>71,72,77</sup> demonstrating that sinoaortic denervation completely blocked autonomic and cardiovascular benefits induced by aerobic training in the SHR.

Besides the autonomic adaptive responses, arterial pressure fall observed in trained hypertensive individuals is also dependent on training-induced structural changes in the peripheral vasculature. Previous data from our group demonstrated that 8-12 weeks of moderate aerobic training normalize the wall/lumen ratio of skeletal muscle, heart and diaphragm arterioles that contributed to the attenuation of total peripheral resistance, thus reducing arterial pressure.<sup>68,73,74</sup> Increased nitric oxide and tetrahydrobiopterin bioavailability and decreased NOS uncoupling, superoxide formation and inducible NOS gene expression, were described as the cellular mechanisms related to vascular benefits of aerobic training in the SHR.<sup>81</sup>

## CONCLUSIONS

Autonomic dysfunction, characterized by sympathovagal unbalance associated with baroreflex dysfunction, contributes widely to the establishment/maintenance of hypertension. Increased oxidative stress and pro-inflammatory profile observed in the SHR contribute to baroreflex dysfunction, augment sympathetic activity and cause vascular and tissue deleterious remodeling. Reduced baroreflex sensitivity implies in increased



**Figure2:** Training-induced time-course adaptive changes on autonomic control and cardiovascular parameters in the SHR. Only 2 weeks of aerobic training (T) were enough to normalize oxidative stress and pro-inflammatory profile within the Paraventricular nucleus of the hypothalamus (PVN) and to correct both cardiac vagal activity and baroreflex dysfunction. These training-induced responses precede the augmentation of heart rate variability and the appearance of resting bradycardia (at the 4<sup>th</sup> week) as well as the partial reduction of peripheral vascular resistance (related to the reversion of arteriolar hypertrophic remodeling in exercised tissues) and a 9% fall in the arterial pressure (both occurring at the 8<sup>th</sup> week of training). Modified from Masson & Michelini, 2014.<sup>64</sup>

arterial pressure variability, sympathetic and RAS activation and the appearance of end-organ damage, which potentiate the development/maintenance of hypertension and constitute a positive feedback mechanism to perpetuate the hypertensive disease.

In the treatment of hypertension, it is crucial to promptly normalize autonomic dysfunction that decreases tissue injuries. It is important to note that aerobic training blocks oxidative stress and inflammation in the autonomic brain areas, normalizes baroreflex function and attenuates others autonomic and cardiovascular dysfunctions related to hypertension, even in the persistence of elevated arterial pressure. Thus, aerobic training constitutes an important therapeutic tool to decrease cardiovascular mortality in hypertensive patients.

#### FINANCIAL SUPPORT

Coordination of Higher Education Personnel Training (CAPES fellowship to GSM), Foundation of the State of São Paulo (FAPESP proc. No. 2011 / 91410-9, research Grant to LCM) and National Research Council (CNPq, fellowship to LCM).

#### REFERENCES

1. Michelini LC. The NTS and Integration of Cardiovascular Control during Exercise in Normotensive and Hypertensive Individuals. *Curr Hypertens Rep.* 2007; 9(3): 214-221. doi: [10.1007/s11906-007-0039-x](https://doi.org/10.1007/s11906-007-0039-x)
2. Kikuya M, Hozawa A, Ohokubo T, et al. Prognostic Significance of Blood Pressure and Heart Rate Variabilities: the OhasamaStudy. *Hypertension.* 2000; 36(5): 901-906. doi: [10.1161/01.HYP.36.5.901](https://doi.org/10.1161/01.HYP.36.5.901)
3. Chobanian AV, Bakris GL, Black HR, et al. National Heart, Lung, and Blood Institute Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure; National High Blood Pressure Education Program Coordinating Committee. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA.* 2003; 289(19): 2560-2572.
4. Minami N, Imai Y, Munakata M, et al. Age-related Changes in Blood Pressure, Heart Rate and Baroreflex Sensitivity in SHR. *Clin Exp Pharmacol Physiol Suppl.* 1898; 15: 85-87. doi: [10.1111/j.1440-1681.1989.tb02999.x](https://doi.org/10.1111/j.1440-1681.1989.tb02999.x)
5. Cates MJ, Steed PW, Abdala AP, et al. Elevated Vertebrobasilar Artery Resistance in Neonatal Spontaneously Hypertensive Rats. *J Appl Physiol.* 2011; 111(1): 149-156. doi: [10.1152/jappphysiol.00220.2011](https://doi.org/10.1152/jappphysiol.00220.2011)
6. Shu HF, Wang BR, Wang SR, et al. IL-1beta Inhibits IK and Increases [Ca<sup>2+</sup>]<sub>i</sub> in the Carotid Body Glomus Cells and Increases Carotid Sinus Nerve Firings in the Rat. *Eur J Neurosci.* 2007; 25(12): 3638-3647. doi: [10.1111/j.1460-9568.2007.05586.x](https://doi.org/10.1111/j.1460-9568.2007.05586.x)
7. McBryde FD, Abdala AP, Hendy EB, et al. The Carotid Body as a Putative Therapeutic Target for the Treatment of Neurogenic Hypertension. *Nat Commun.* 2013; 4: 2395. doi: [10.1038/ncomms3395](https://doi.org/10.1038/ncomms3395)
8. Kang, YM, Ma Y, Zheng JP, et al. Brain Nuclear Factor-Kappa B Activation Contributes to Neurohumoral Excitation in Angiotensin II-Induced Hypertension. *Cardiovasc Res.* 2009; 82(3): 503-512. doi: [10.1093/cvr/cvp073](https://doi.org/10.1093/cvr/cvp073)
9. Sinnayah P, Lazartigues E, Sakai K, Sharma RV, Sigmund CD, Davisson RL. Genetic Ablation of Angiotensinogen in the Subfornical Organ of the Brain Prevents the Central Angiotensinergic Pressor Response. *Circ Res.* 2006; 99(10): 1125-1131. doi: [10.1161/01.RES.0000250259.66683.f5](https://doi.org/10.1161/01.RES.0000250259.66683.f5)
10. Biancardi VC, Son SJ, Ahmadi S, Filosa JA, Stern JE. Circulating Angiotensin II Gains Access to the Hypothalamus and Brain Stem During Hypertension via Breakdown of the Blood-Brain Barrier. *Hypertension.* 2014; 63(3): 572-579. doi: [10.1161/HYPERTENSIONAHA.113.01743](https://doi.org/10.1161/HYPERTENSIONAHA.113.01743)
11. Waki H, Bhuiyan ME, Gouraud SS, et al. Acute Reductions in Blood Flow Restricted to the Dorsomedial Medulla Induce a Pressor Response in Rats. *J Hypertens.* 2011; 29(8): 1536-1545. doi: [10.1097/HJH.0b013e3283484106](https://doi.org/10.1097/HJH.0b013e3283484106)
12. Fan J, Zhang B, Shu HF, et al. Interleukin-6 Increases Intracellular Ca<sup>2+</sup> Concentration and Induces Catecholamine Secretion in Rat Carotid Body Glomus Cells. *J Neurosci Res.* 2009; 87(12): 2757-2762. doi: [10.1002/jnr.2210](https://doi.org/10.1002/jnr.2210)
13. Chae CU, Lee RT, Rifai N, Ridker PM. Blood Pressure and Inflammation in Apparently Healthy Men. *Hypertension.* 2001; 38(3): 399-403. doi: [10.1161/01.HYP.38.3.399](https://doi.org/10.1161/01.HYP.38.3.399)
14. Haensel A, Mills PJ, Nelesen RA, Ziegler MG, Dimsdale JE. The Relationship Between Heart Rate Variability and Inflammatory Markers in Cardiovascular Diseases. *Psychoneuroendocrinology.* 2008; 33(10): 1305-1312. doi: [10.1016/j.psyneuen.2008.08.007](https://doi.org/10.1016/j.psyneuen.2008.08.007)
15. Mkrtchian S, Kählin J, Ebberyd A, et al. The Human Carotid Body Transcriptome with Focus on Oxygen Sensing and Inflammation—a Comparative Analysis. *J Physiol.* 2012; 590: 3807-3819. doi: [10.1113/jphysiol.2012.231084](https://doi.org/10.1113/jphysiol.2012.231084)
16. Sakai K, Agassandian K, Morimoto S, et al. Local Production of Angiotensin II in the Subfornical Organ Causes Elevated Drinking. *J Clin Invest.* 2007; 117(4): 1088-1095. doi: [10.1172/JCI1242](https://doi.org/10.1172/JCI1242)
17. Oliveira-Sales EB, Colombari DS, Davisson RL, et al. Kidney-Induced Hypertension Depends on Superoxide Signaling in

- the Rostral Ventrolateral Medulla. *Hypertension*. 2010; 56(2): 290-296. doi: [10.1161/HYPERTENSIONAHA.110.150425](https://doi.org/10.1161/HYPERTENSIONAHA.110.150425)
18. Yamazato M, Yamazato Y, Sun C, Diez-Freire C, Raizada MK. Overexpression of Angiotensin-Converting Enzyme 2 in the Rostral Ventrolateral Medulla Causes Long-Term Decrease in Blood Pressure in the Spontaneously Hypertensive Rats. *Hypertension*. 2007; 49(4): 926-931. doi: [10.1161/01.HYP.0000259942.38108.20](https://doi.org/10.1161/01.HYP.0000259942.38108.20)
19. Xu H, Oliveira-Sales EB, McBride F, et al. Upregulation of Junctional Adhesion Molecule-A is a Putative Prognostic Marker of Hypertension. *Cardiovasc Res*. 2012; 96(3): 552-560. doi: [10.1093/cvr/cvs273](https://doi.org/10.1093/cvr/cvs273)
20. Coleman CG, Wang G, Faraco G, et al. Membrane Trafficking of NADPH Oxidase p47 (phox) in Paraventricular Hypothalamic Neurons Parallels Local Free Radical Production in Angiotensin II Slow-pressor Hypertension. *J Neurosci*. 2013; 33(10): 4308-4316. doi: [10.1523/JNEUROSCI.3061-12.2013](https://doi.org/10.1523/JNEUROSCI.3061-12.2013)
21. Wang G, Sarkar P, Peterson JR, et al. COX-1-derived PGE2 and PGE2 Type 1 Receptors Are Vital for Angiotensin II-Induced Formation of Reactive Oxygen Species and Ca(2+) Influx in the Subfornical Organ. *Am J Physiol Heart Circ Physiol*. 2013; 305(10): H1451-H1461. doi: [10.1152/ajpheart.00238.2013](https://doi.org/10.1152/ajpheart.00238.2013)
22. Chan SH, Wang LL, Tseng HL, Chan JY. Upregulation of AT<sub>1</sub> Receptor Gene on Activation of Protein Kinase Cbeta/nicotinamide Adenine Dinucleotide Diphosphate Oxidase/ERK1/2/c-fos Signaling Cascade Mediates Long-term Pressor Effect of Angiotensin II in Rostral Ventrolateral Medulla. *J Hypertens*. 2007; 25(9): 1845-1861.
23. Wang G, Anrather J, Glass MJ, et al. Nox2, Ca<sup>2+</sup>, and Protein Kinase C Play a Role in Angiotensin II-Induced Free Radical Production in Nucleus Tractus Solitarius. *Hypertension*. 2006; 48(3): 482-489. doi: [10.1161/01.HYP.0000236647.55200.07](https://doi.org/10.1161/01.HYP.0000236647.55200.07)
24. Yu Y, Xue BJ, Zhang ZH, et al. Early Interference with p44/42 Mitogen-Activated Protein Kinase Signaling in Hypothalamic Paraventricular Nucleus Attenuates Angiotensin II-Induced Hypertension. *Hypertension*. 2013; 61(4): 842-849. doi: [10.1161/HYPERTENSIONAHA.111.00080](https://doi.org/10.1161/HYPERTENSIONAHA.111.00080)
25. Cardinale JP, Sriramula S, Mariappan N, Agarwal D, Francis J. Angiotensin II-Induced Hypertension is Modulated by Nuclear Factor-κB in the Paraventricular Nucleus. *Hypertension*. 2012; 59(1): 113-121. doi: [10.1161/HYPERTENSIONAHA.111.182154](https://doi.org/10.1161/HYPERTENSIONAHA.111.182154)
26. Wang G, Coleman CG, Glass MJ, et al. Angiotensin II Type 2 Receptor-Coupled Nitric Oxide Production Modulates Free Radical Availability and Voltage-gated Ca<sup>2+</sup> Currents in NTS Neurons. *Am J Physiol Regul Integr Comp Physiol*. 2012; 302(9): R1076-R1083. doi: [10.1152/ajpregu.00571.2011](https://doi.org/10.1152/ajpregu.00571.2011)
27. Nakamura T, Lipton SA. Redox Modulation by S-nitrosylation Contributes to Protein Misfolding, Mitochondrial Dynamics, and Neuronal Synaptic Damage in Neurodegenerative Diseases. *Cell Death Differ*. 2011; 18(9):1478-1486. doi: [10.1038/cdd.2011.65](https://doi.org/10.1038/cdd.2011.65)
28. Lob HE, Schultz D, Marvar PJ, Davisson RL, Harrison DG. Role of the NADPH Oxidases in the Subfornical Organ in Angiotensin II-Induced Hypertension. *Hypertension*. 2013; 61(2): 382-387. doi: [10.1161/HYPERTENSIONAHA.111.00546](https://doi.org/10.1161/HYPERTENSIONAHA.111.00546)
29. Cao X, Peterson JR, Wang G, et al. Angiotensin II-Dependent Hypertension Requires Cyclooxygenase 1-Derived Prostaglandin E2 and EP1 Receptor Signaling in the Subfornical Organ of the Brain. *Hypertension*. 2012; 59(4): 869-876. doi: [10.1161/HYPERTENSIONAHA.111.182071](https://doi.org/10.1161/HYPERTENSIONAHA.111.182071)
30. Zimmerman MC, Dunlay RP, Lazartigues E, et al. Requirement for Rac1-Dependent NADPH Oxidase in the Cardiovascular and Dipsogenic Actions of Angiotensin II in the Brain. *Circ Res*. 2004; 95(5): 532-539. doi: [10.1161/01.RES.0000139957.22530.b9](https://doi.org/10.1161/01.RES.0000139957.22530.b9)
31. Burmeister MA, Young CN, Braga VA, Butler SD, Sharma RV, Davisson RL. In Vivo Bioluminescence Imaging Reveals Redox-Regulated Activator Protein-1 Activation in Paraventricular Nucleus of Mice with Renovascular Hypertension. *Hypertension*. 2011; 57(2): 289-297. doi: [10.1161/HYPERTENSIONAHA.110.160564](https://doi.org/10.1161/HYPERTENSIONAHA.110.160564)
32. Oliveira-Sales EB, Nishi EE, Carillo BA, et al. Oxidative Stress in the Sympathetic Premotor Neurons Contributes to Sympathetic Activation in Renovascular Hypertension. *Am J Hypertens*. 2009; 22(5): 484-492. doi: [10.1038/ajh.2009.17](https://doi.org/10.1038/ajh.2009.17)
33. Oliveira-Sales EB, Dugaich AP, Carillo BA, et al. Oxidative Stress Contributes to Renovascular Hypertension. *Am J Hypertens*. 2008; 21(1): 98-104. doi: [10.1038/ajh.2007.12](https://doi.org/10.1038/ajh.2007.12)
34. Nishihara M, Hirooka Y, Matsukawa R, Kishi T, Sunagawa K. Oxidative Stress in the Rostral Ventrolateral Medulla Modulates Excitatory and Inhibitory Inputs in Spontaneously Hypertensive Rats. *J Hypertens*. 2012; 30(1): 97-106. doi: [10.1097/HJH.0b013e32834e1df4](https://doi.org/10.1097/HJH.0b013e32834e1df4)
35. Tai MH, Wang LL, Wu KL, Chan JY. Increased Superoxide Anion in Rostral Ventrolateral Medulla Contributes to Hypertension in Spontaneously Hypertensive Rats via Interactions with Nitric Oxide. *Free Radic Biol Med*. 2005; 38(4): 450-462. doi: [10.1016/j.freeradbiomed.2004.11.015](https://doi.org/10.1016/j.freeradbiomed.2004.11.015)
36. Kishi T, Hirooka Y, Kimura Y, Ito K, Shimokawa H, Takeshita A. Increased Reactive Oxygen Species in Rostral Ventrolateral Medulla Contribute to Neural Mechanisms of Hypertension in Stroke-Prone Spontaneously Hypertensive Rats. *Circulation*. 2004; 109(19): 2357-2362. doi: [10.1161/01](https://doi.org/10.1161/01)

CIR.0000128695.49900.12

37. Tsai CY, Su CH, Baudrie V, et al. Visualizing Oxidative Stress-Induced Depression of Cardiac Vagal Baroreflex by MRI/DTI in a Mouse Neurogenic Hypertension Model. *NeuroImage*. 2013; 82: 190-199. doi: [10.1016/j.neuroimage.2013.05.124](https://doi.org/10.1016/j.neuroimage.2013.05.124)
38. Wei SG, Zhang ZH, Beltz TG, Yu Y, Johnson AK, Felder RB. Subfornical Organ Mediates Sympathetic and Hemodynamic Responses to Blood-borne Proinflammatory Cytokines. *Hypertension*. 2013; 62(1): 118-125. doi: [10.1161/HYPERTENSIONAHA.113.01404](https://doi.org/10.1161/HYPERTENSIONAHA.113.01404)
39. Shi P, Diez-Freire C, Jun JY, et al. Brain Microglial Cytokines in Neurogenic Hypertension. *Hypertension*. 2010; 56(2): 297-303. doi: [10.1161/HYPERTENSIONAHA.110.150409](https://doi.org/10.1161/HYPERTENSIONAHA.110.150409)
40. Takagishi M, Waki H, Bhuiyan ME, et al. IL-6 Microinjected in the Nucleus Tractus Solitarius Attenuates Cardiac Baroreceptor Reflex Function in Rats. *Am J Physiol Regul Integr Comp Physiol*. 2010; 298(1): R183-R190. doi: [10.1152/ajpregu.00176.2009](https://doi.org/10.1152/ajpregu.00176.2009)
41. Amann FW, Bolli P, Kiowski W, Bühler FR. Enhanced Alpha-Adrenoreceptor-Mediated Vasoconstriction in Essential Hypertension. *Hypertension*. 1981; 3(2): I119-I123. doi: [10.1161/01.HYP.3.3.Pt\\_2.I119](https://doi.org/10.1161/01.HYP.3.3.Pt_2.I119)
42. Rokosh DG, Simpson PC. Knockout of the Alpha 1A/C-Adrenergic Receptor Subtype: the Alpha 1A/C is expressed in Resistance Arteries and is required to Maintain Arterial Blood Pressure. *Proc Natl Acad Sci USA*. 2002; 99(14): 9474-9479. doi: [10.1073/pnas.132552699](https://doi.org/10.1073/pnas.132552699)
43. Erami C, Zhang H, Tanoue A, Tsujimoto G, Thomas SA, Faber JE. Adrenergic Catecholamine Trophic Activity Contributes to Flow-Mediated Arterial Remodeling. *Am J Physiol Heart Circ Physiol*. 2005; 289(2): H744-H753. doi: [10.1152/ajpheart.00129.2005](https://doi.org/10.1152/ajpheart.00129.2005)
44. Gayen JR, Zhang K, Rao RSP, et al. Role of Reactive Oxygen Species in Hyperadrenergic Hypertension: Biochemical, Physiological, and Pharmacological Evidence from Targeted Ablation of the Chromogranin a (Chga) Gene. *Circ Cardiovasc Genet*. 2010; 3(5): 414-425. doi: [10.1161/CIRCGENETICS.109.924050](https://doi.org/10.1161/CIRCGENETICS.109.924050)
45. Burgi K, Cavalleri MT, Alves AS, Britto LR, Antunes VR, Michelini LC. Tyrosine Hydroxylase Immunoreactivity as Indicator of Sympathetic Activity: Simultaneous Evaluation in Different Tissues of Hypertensive Rats. *Am J Physiol Regul Integr Comp Physiol*. 2011; 300(2): R264-R271. doi: [10.1152/ajpregu.00687.2009](https://doi.org/10.1152/ajpregu.00687.2009)
46. Osborn JW, Fink GD. Region-Specific Changes in Sympathetic Nerve Activity in Angiotensin II-Salt Hypertension in the Rat. *Exp Physiol*. 2010; 95(1): 61-68. doi: [10.1113/expphysiol.2008.046326](https://doi.org/10.1113/expphysiol.2008.046326)
47. Masson S, Arosio B, Luvarà G, et al. Remodelling of Cardiac Extracellular Matrix During Beta-Adrenergic Stimulation: Upregulation of SPARC in the Myocardium of Adult Rats. *J Mol Cell Cardiol*. 1998; 30(8): 1505-1514. doi: [10.1006/jmcc.1998.0714](https://doi.org/10.1006/jmcc.1998.0714)
48. Briest W, Homagk L, Rassler B, et al. Norepinephrine-Induced Changes in Cardiac Transforming Growth Factor-Beta Isoform Expression Pattern of Female and Male Rats. *Hypertension*. 2004; 44(4): 410-418. doi: [10.1161/01.HYP.0000141414.87026.4d](https://doi.org/10.1161/01.HYP.0000141414.87026.4d)
49. Levick SP, Murray DB, Janicki JS, Brower GL. Sympathetic Nervous System Modulation of Inflammation and Remodeling in the Hypertensive Heart. *Hypertension*. 2010; 55(2): 270-276. doi: [10.1161/HYPERTENSIONAHA.109.142042](https://doi.org/10.1161/HYPERTENSIONAHA.109.142042)
50. Schlaich MP, Kaye DM, Lambert E, Sommerville M, Socratous F, Esler MD. Relation between Cardiac Sympathetic Activity and Hypertensive Left Ventricular Hypertrophy. *Circulation*. 2003; 108(5): 560-565. doi: [10.1161/01.CIR.0000081775.72651.B6](https://doi.org/10.1161/01.CIR.0000081775.72651.B6)
51. Elhawary AM, Pang CC. Alpha 1b-Adrenoceptors Mediate Renal Tubular Sodium and Water Reabsorption in the Rat. *Br J Pharmacol*. 1994; 111(3): 819-824.
52. Hesse IF, Johns EJ. The Role of Alpha-Adrenoceptors in the Regulation of Renal Tubular Sodium Reabsorption and Renin Secretion in the Rabbit. *Br J Pharmacol*. 1985; 84(3): 715-724.
53. Graham RM, Pettinger WA, Sagalowsky A, Brabson J, Gandler T. Renal Alpha-Adrenergic Receptor Abnormality in the Spontaneously Hypertensive Rat. *Hypertension*. 1982; 4(6): 881-887. doi: [10.1161/01.HYP.4.6.881](https://doi.org/10.1161/01.HYP.4.6.881)
54. Amann K, Koch A, Hofstetter J, et al. Glomerulosclerosis and Progression: Effect of Subantihypertensive Doses of Alpha and Beta Blockers. *Kidney Int*. 2001; 60(4): 1309-1323. doi: [10.1046/j.1523-1755.2001.00936.x](https://doi.org/10.1046/j.1523-1755.2001.00936.x)
55. Nakaya H, Sasamura H, Kitamura Y, et al. Effects of Angiotensin Inhibitors on Renal Injury and Angiotensin Receptor Expression in Early Hypertensive Nephrosclerosis. *Hypertens Res*. 1999; 22(4): 303-312. doi: [10.1291/hypres.22.303](https://doi.org/10.1291/hypres.22.303)
56. Higashikuni Y, Takaoka M, Iwata H, et al. Aliskiren in Combination with Valsartan Exerts Synergistic Protective Effects Against Ventricular Remodeling After Myocardial Infarction in Mice. *Hypertens Res*. 2012; 35(1): 62-69. doi: [10.1038/hr.2011.136](https://doi.org/10.1038/hr.2011.136)
57. Yamamoto E, Kataoka K, Dong YF, et al. Aliskiren Enhances the Protective Effects of Valsartan Against Cardiovascular and Renal Injury in Endothelial Nitric Oxide Synthase-Deficient Mice. *Hypertension*. 2009; 54(3): 633-638. doi: [10.1161/HYPERTENSIONAHA.109.133884](https://doi.org/10.1161/HYPERTENSIONAHA.109.133884)



58. Zhuo JL, Li XC. New Insights and Perspectives on Intra-renal Renin-Angiotensin System: Focus on Intracrine/Intracellular Angiotensin II. *Peptides*. 2011; 32(7): 1551-1565. doi: [10.1016/j.peptides.2011.05.012](https://doi.org/10.1016/j.peptides.2011.05.012)
59. Li XC, Zhuo JL. Proximal Tubule-Dominant Transfer of AT (1a) Receptors Induces Blood Pressure Responses to Intracellular Angiotensin II in AT (1a) Receptor-Deficient Mice. *Am J Physiol Regul Integr Comp Physiol*. 2013; 304(8): R588-R598. doi: [10.1152/ajpregu.00338.2012](https://doi.org/10.1152/ajpregu.00338.2012)
60. Nosaka S, Murata K, Kobayashi M, Cheng ZB, Maruyama J. Inhibition of Baroreflex Vagal Bradycardia by Activation of the Rostral Ventrolateral Medulla in Rats. *Am J Physiol Heart Circ Physiol*. 2000; 279(3): H1239-H1247.
61. Ormezzano O, Cracowski JL, Quesada JL, Pierre H, Mallion JM, Baguet JP. Evaluation of the Prognostic Value of Baroreflex Sensitivity in Hypertensive Patients: the EVABAR Study. *J Hypertens*. 2008; 26(7): 1373-1378. doi: [10.1097/HJH.0b013e3283015e5a](https://doi.org/10.1097/HJH.0b013e3283015e5a)
62. Yasuoka S, Kai H, Kajimoto H, et al. Blood Pressure Variability Activates Cardiac Mineralocorticoid Receptor and Induces Cardiac Remodeling in Hypertensive Rats. *Circ J*. 2013; 77(6): 1474-1481. doi: [10.1253/circj.CJ-12-1253](https://doi.org/10.1253/circj.CJ-12-1253)
63. Kudo H, Kai H, Kajimoto H, et al. Exaggerated Blood Pressure Variability Superimposed on Hypertension Aggravates Cardiac Remodeling in Rats via Angiotensin II System-Mediated Chronic Inflammation. *Hypertension*. 2009; 54(4): 832-838. doi: [10.1161/HYPERTENSIONAHA.109.135905](https://doi.org/10.1161/HYPERTENSIONAHA.109.135905)
64. Masson GS, Michelini LC. Sympathetic Nervous Activity and Development Organ-Alvona injuries Hypertension: Autonomic Benefits Induced Aerobic Training. *Rev Soc Bras Cardiol Est Sao Paulo*. 2014; 2(4): 16-24.
65. Cornelissen VA, Smart NA. Exercise Training for Blood Pressure: a Systematic Review and Meta-Analysis. *J Am Heart Assoc*. 2013; 2(1): e004473. doi: [10.1161/JAHA.112.004473](https://doi.org/10.1161/JAHA.112.004473)
66. Agarwal D, Elks CM, Reed SD, Mariappan N, Majid DS, Francis J. Chronic Exercise Preserves Renal Structure and Hemodynamics in Spontaneously Hypertensive Rats. *Antioxid Redox Signal*. 2012; 16(2): 139-152. doi: [10.1089/ars.2011.396](https://doi.org/10.1089/ars.2011.396)
67. Rossoni LV, Oliveira RA, Caffaro RR, et al. Cardiac Benefits of Exercise Training in Aging Spontaneously Hypertensive Rats. *J Hypertens*. 2011; 29(12): 2349-2358. doi: [10.1097/HJH.0b013e32834d2532](https://doi.org/10.1097/HJH.0b013e32834d2532)
68. Amaral SL, Silveira NP, Zorn TM, Michelini LC. Exercise Training Causes Skeletal Muscle Venular Growth and Alters Hemodynamic Responses in Spontaneously Hypertensive Rats. *J Hypertens*. 2001; 19(5): 931-940.
69. Laterza MC, De Matos LD, Trombetta IC, et al. Exercise Training Restores Baroreflex Sensitivity in Never-Treated Hypertensive Patients. *Hypertension*. 2007; 49(6): 1298-1306. doi: [10.1161/HYPERTENSIONAHA.106.085548](https://doi.org/10.1161/HYPERTENSIONAHA.106.085548)
70. Masson GS, Costa Ts, Yshii L, et al. Time-Dependent Effects of Training on Cardiovascular Control in Spontaneously Hypertensive Rats: Role for Brain Oxidative Stress and Inflammation and Baroreflex Sensitivity. *PLoS One*. 2014, 9(5): e94927. doi: [10.1371/journal.pone.0094927](https://doi.org/10.1371/journal.pone.0094927)
71. Ceroni A, Chaar LJ, Bombein RL, Michelini LC. Chronic Absence of Baroreceptor Inputs Prevents Training-Induced Cardiovascular Adjustments in Normotensive and Spontaneously Hypertensive Rats. *Exp Physiol*. 2009; 94(6): 630-640. doi: [10.1113/expphysiol.2008.046128](https://doi.org/10.1113/expphysiol.2008.046128)
72. Moraes-Silva IC, De La Fuente RN, Mostarda C, et al. Baroreflex Deficit Blunts Exercise Training-Induced Cardiovascular and Autonomic Adaptations in Hypertensive Rats. *Clin Exp Pharmacol Physiol*. 2010; 37(3): e114-e120. doi: [10.1111/j.1440-1681.2009.05333.x](https://doi.org/10.1111/j.1440-1681.2009.05333.x)
73. Melo RM, Martinho E Jr, Michelini LC. Training-Induced, Pressure-Lowering Effect in SHR: Wide Effects on Circulatory Profile of Exercised and Nonexercised Muscles. *Hypertension*. 2003; 42(4): 851-857. doi: [10.1161/01.HYP.0000086201.27420.33](https://doi.org/10.1161/01.HYP.0000086201.27420.33)
74. Brum PC, Da Silva GJ, Moreira ED, Ida F, Negrão CE, Krieger EM. Exercise Training Increases Baroreceptor Gain Sensitivity in Normal and Hypertensive Rats. *Hypertension*. 2000; 36(6): 1018-1022. doi: [10.1161/01.HYP.36.6.1018](https://doi.org/10.1161/01.HYP.36.6.1018)
75. Stern JE, Sonner PM, Son SJ, Silva FC, Jackson K, Michelini LC. Exercise Training Normalizes an Increased Neuronal Excitability of NTS-Projecting Neurons of the Hypothalamic Paraventricular Nucleus in Hypertensive Rats. *J Neurophysiol*. 2012; 107(10): 2912-2921. doi: [10.1152/jn.00884.2011](https://doi.org/10.1152/jn.00884.2011)
76. Michelini LC, Stern JE. Exercise-Induced Neuronal Plasticity in Central Autonomic Networks: Role in Cardiovascular Control. *Exp Physiol*. 2009; 94(9): 947-960. doi: [10.1113/expphysiol.2009.047449](https://doi.org/10.1113/expphysiol.2009.047449)
77. Cavalleri MT, Burgi K, Cruz JC, Jordão MT, Ceroni A, Michelini LC. Afferent Signaling Drives Oxytocinergic Preautonomic Neurons and Mediates Training-Induced Plasticity. *Am J Physiol Regul Integr Comp Physiol*. 2011; 301(4): R958-R966. doi: [10.1152/ajpregu.00104.2011](https://doi.org/10.1152/ajpregu.00104.2011)
78. Higa-Taniguchi KT, Felix JV, Michelini LC. Brainstem Oxytocinergic Modulation of Heart Rate Control in Rats: Effects of Hypertension and Exercise Training. *Exp Physiol*. 2009; 94(11): 1103-1113. doi: [10.1113/expphysiol.2009.049262](https://doi.org/10.1113/expphysiol.2009.049262)

79. Heusser K, Tank J, Engeli S, et al. Carotid Baroreceptor Stimulation, Sympathetic Activity, Baroreflex Function, and Blood Pressure in Hypertensive Patients. *Hypertension*. 2010; 55(3): 619-626. doi: [10.1161/HYPERTENSIONAHA.109.140665](https://doi.org/10.1161/HYPERTENSIONAHA.109.140665)

80. Wustmann K, Kucera JP, Scheffers I, et al. Effects of Chronic Baroreceptor Stimulation on the Autonomic Cardiovascular Regulation in Patients with Drug-Resistant Arterial Hypertension. *Hypertension*. 2009; 54(3): 530-536. doi: [10.1161/HYPERTENSIONAHA.109.134023](https://doi.org/10.1161/HYPERTENSIONAHA.109.134023)

81. Roque FR, Briones AM, García-Redondo AB, et al. Aerobic Exercise Reduces Oxidative Stress and Improves Vascular Changes of Small Mesenteric and Coronary Arteries in Hypertension. *Br J Pharmacol*. 2013; 168(3): 686-703.